

Quark masses using domain wall fermions *

Matthew Wingate^a

^aRIKEN BNL Research Center, Brookhaven National Laboratory, Upton, NY 11973, USA

Due to the attractive features that domain wall fermions possess with respect to chiral symmetry, we continue our investigation of the light quark masses with this discretization. Achieving reliable results, especially for $(m_u + m_d)/2$, requires strict control of systematic uncertainties. Our present results were obtained on a quenched $\beta = 6.0$ lattice with spatial volume $\approx (1.5 \text{ fm})^3$. Consequently we remark on effects of finite volume as well as finite extent in the fictitious fifth dimension. We compute the renormalization factors nonperturbatively and compare to the one-loop perturbative result.

1. INTRODUCTION

Last year an exploratory calculation of the strange quark mass was completed using the domain wall fermion discretization within the quenched approximation [1]. At the 15–20% level the results were encouraging: the pion mass squared extrapolated to zero at the input mass $m \rightarrow 0$, simulations at three different lattice spacings gave similar values for $m_s^{\overline{\text{MS}}}(2 \text{ GeV})$ and the values were in agreement with other lattice results. This work reports on the progress of the RIKEN/BNL/CU (RBC) Collaboration toward a precise calculation of the strange quark mass.

2. SPECTRUM

Using domain wall quarks we have computed light hadron masses on a $16^3 \times 32$ quenched lattice with the following parameters: $\beta = 6.0$ (plaquette action), domain wall height $M = 1.8$, seven values of valence quark masses (am 's), and number of sites in the extra dimension $N_s = 16$. Results from 85 configurations for the pseudoscalar meson, vector meson, and nucleon masses are shown in Figures 1 and 2 (see also Ref. [2]).

With the light masses and improved statistics of this study compared to Ref. [1], it has become clear that the linear extrapolation of $(aM_\pi)^2$ to $am = 0$ gives a positive intercept, $0.018(2)$ (sta-

tistical error only), for these simulation parameters (see Fig. 1). Several effects could be responsible for this feature. First, the finite (and relatively small) volume of our lattice could increase the mass of the pion at a given am , and these effects would be especially visible at the lighter masses; as shown below the smallest am , 0.01, is roughly 1/4 the strange quark mass. Second, there is an intrinsic breaking of chiral symmetry due to the finite N_s : the right- and left-handed surfaces states have some overlap within the fifth dimension and would result in a residual quark mass m_{res} . Both finite V and N_s effects should be mostly am -independent. On the other hand studies by Columbia [3] at $\beta = 5.7$ suggest that in the large N_s limit one cannot account for the entire M_π^2 intercept by increasing the spatial volume by a factor 2^3 . Quenching effects predicted by quenched chiral perturbation theory or due to artificial instantons could play a role. The issue of residual mass is addressed in Ref. [3].

We cannot presently distinguish am -dependent finite V and N_s effects from any nonleading or logarithmic behavior of the pseudoscalar meson mass. For this analysis we assume that the above effects are all am -independent, so then the chiral limit is at $am = -0.0038$. We discuss systematic uncertainties regarding this assumption below.

The am corresponding to the light quark mass is determined by extrapolating $(M_\pi/M_\rho)^2$ to its physical ratio. There $M_N/M_\rho = 1.37(5)$ compared to the physical ratio 1.22. The lattice spacing is then set using the ρ mass, and the strange quark mass is set using either the physical K or ϕ

*Talk given at Lattice '99, Pisa, Italy; work done in collaboration with T. Blum, P. Chen, N. Christ, M. Creutz, C. Dawson, G. Fleming, R. Mawhinney, S. Ohta, S. Sasaki, G. Siegert, A. Soni, P. Vranas, L. Wu, and Y. Zhestkov (RIKEN/BNL/CU Collaboration)




Figure 1. Pseudoscalar meson mass squared vs. am .




Figure 2. Vector meson and nucleon masses vs. am .

mass. The bare quark masses ($m_q = am + 0.0038$) are thus (with statistical errors only):

bare mass	$1/a = 1.91(0.04)$ GeV	
	lattice units	MeV
m_l	0.00166(0.00005)	3.17(0.11)
$m_s(K)$	0.042(0.003)	80(6)
$m_s(\phi)$	0.053(0.004)	101(7)

The large difference in m_s due to whether the K or ϕ meson is used to set the scale is a common feature to quenched lattice simulations. This is succinctly expressed by the parameter $J \equiv M_{K^*}(dM_V/dM_{PS}^2)$ [4]. The value obtained using physical meson masses is 0.48 ± 0.02 (the spread is due to the fact that $(M_\phi - M_{K^*})/(M_{K^*} - M_\rho) \neq 1$); while our result is $J = 0.34(7)$, comparable to calculations with Wilson fermions. In light of this difference, we do not average $m_s(K)$ and $m_s(\phi)$ but quote results for each input separately. The calculation of the ϕ mass in quenched QCD is sub-

ject to systematic error since disconnected graphs are being neglected, and so in the rest of this work we consider only $m_s(K)$.

So far we have not completed an exhaustive study of the systematic errors at this lattice spacing. Therefore, for the present time we give rough estimates. Since M_N/M_ρ is 10% higher than the physical value, our determination of the lattice spacing has an inherent uncertainty of at least 10%. In this work we have assumed that finite V and N_s effects gave a positive m -independent contribution to the pion mass. If we instead assume such effects account for only half of the nonzero intercept [3] then the quark masses above would all decrease by 4 MeV. This is only a 5% change to the strange quark mass, but clearly the extrapolation to the average up and down quark mass is untrustworthy. A better understanding of the systematic uncertainties is necessary before a reliable calculation of m_l can be completed. The bare strange mass from this work is $m_s(\beta = 6.0) = 80 \pm 6$ (stat.) ± 10 (sys.) MeV.

3. NONPERTURBATIVE RENORMALIZATION

The mass renormalization constant has been computed perturbatively to one-loop for domain wall fermions [1,5]. In this work we compute the renormalization constant nonperturbatively using the regularization independent momentum subtraction (RI/MOM) scheme as advocated by the Rome-Southampton group [6].

To determine the renormalization constant for an operator O , we impose the following renormalization condition:

$$\frac{Z_O(\mu a)}{Z_q(\mu a)} \text{Tr} \left[P_O \Lambda_O(pa) \right] \Big|_{p^2=\mu^2} = 1, \quad (1)$$

where P_O projects out the tree-level spin structure of O , Λ_O is the amputated Greens function of O in momentum space, and Z_q is the quark wavefunction renormalization. For fermion bilinear operators $O_\Gamma(x) = \bar{q}(x)\Gamma q(x)$ only a single momentum space propagator is needed to compute the amputated Greens function

$$\Lambda_\Gamma(p) = S^{-1}(p) \langle S(p) \Gamma (\gamma_5 S^\dagger(p) \gamma_5) \rangle S^{-1}(p) \quad (2)$$

where $S(p) = \sum_y \exp(-ipy) \langle q(y) \bar{q}(0) \rangle$. In this work we compute momentum space propagators on 52 configurations [7]. Eqns. (1) and (2) together give Z_Γ/Z_q at any momentum p . In order for the RI/MOM method to work reliably the momentum should satisfy $\Lambda_{\text{QCD}} \ll p \ll 1/a$ so that nonperturbative and discretization effects, respectively, are small; however with presently accessible lattices this range is narrow. The approach which we follow is to fit to the leading $(ap)^2$ errors [8]. First, we compute Z_q through

$$Z'_q \equiv -\frac{i}{12} \frac{\text{Tr} \sum_\mu \gamma_\mu p_\mu S^{-1}}{4 \sum_\mu p_\mu^2} \quad (3)$$

which differs from Z_q at $O(g^4)$ in Landau gauge [9], then multiply Z_Γ/Z_q by Z'_q to give Z'_Γ and divide by the two-loop running c'_Γ [9] resulting in

$$Z_\Gamma^{\text{RGI}}(a) = \frac{Z'_\Gamma(\mu a)}{c'_\Gamma(\mu)} = \frac{Z_\Gamma^{\overline{\text{MS}}}(\mu a)}{c_\Gamma^{\overline{\text{MS}}}(\mu)} \quad (4)$$

which is scale independent through two-loop order. The result for the scalar density is shown in Fig. 3. The solid line is a linear fit to $O(a^2 p^2)$ discretization errors, and we take the extrapolated value at $(ap)^2 = 0$ as our Z_S^{RGI} . $Z_S^{\overline{\text{MS}}}$ at 2 GeV is obtained by multiplying by the two-loop running factor $c_S^{\overline{\text{MS}}}(2 \text{ GeV})$. The result for $Z_m^{\overline{\text{MS}}} \equiv 1/Z_S^{\overline{\text{MS}}}$ is $1.63(7)(9)$ (stat.)(sys.) which should be compared to 1.32, the one-loop perturbative Z_m [1,5] for these simulation parameters.

Taking the lattice $m_s(K)$ from the previous section we find at $\beta = 6.0$

$$m_s^{\overline{\text{MS}}}(2 \text{ GeV}) = 130 \pm 11 \pm 18 \text{ MeV} \quad (5)$$

where the first error is the statistical uncertainty and the second is the systematic uncertainty.

4. CONCLUSIONS

We have completed a calculation of the light and strange quark masses at one lattice spacing and volume within the quenched lattice spacing. The increased precision has exposed finite N_s and V effects and these lead to errors of roughly 5% in the strange quark mass. We will soon have a calculation of m_s on a coarser ($\beta = 5.85$) lattice, and of course further work at larger volumes,




Figure 3. Scalar density renormalization factor with perturbative scale dependence removed up to $O(g^6)$. The solid line is a linear fit to $a^2 p^2$ discretization effects.

larger N_s , and smaller lattice spacing will reduce systematic uncertainties and yield a precise determination of the strange quark mass.

ACKNOWLEDGMENTS

The RBC Collaboration is grateful to RIKEN, Brookhaven National Laboratory, and the U.S. Department of Energy for providing the facilities essential for the completion of this work.

REFERENCES

1. T. Blum, A. Soni, M. Wingate, Nucl. Phys. B (Proc. Suppl.) **73**, 201 (1999); *ibid.*, to appear in Phys. Rev. D, hep-lat/9902016.
2. L. Wu, these proceedings.
3. G. Fleming, these proceedings.
4. P. Lacey and C. Michael, Phys. Rev. **D52**, 5213 (1995).
5. S. Aoki, T. Izubuchi, Y. Kuramashi, and Y. Taniguchi, Phys. Rev. **D59**, 094505 (1999).
6. G. Martinelli *et al.*, Nucl. Phys. **B445**, 81 (1995).
7. C. Dawson, these proceedings.
8. V. Giménez *et al.*, Nucl. Phys. **B531**, 429 (1998).
9. E. Franco and V. Lubicz, Nucl. Phys. **B531**, 641 (1998).